

REJUVENATION RESEARCH
Volume 11, Number 2, 2008

<http://www.ncbi.nlm.nih.gov/pubmed/18321197>

<http://www.liebertonline.com/doi/abs/10.1089/rej.2008.0661>

Scientific Justification of Cryonics Practice

Benjamin P. Best*

ABSTRACT

Very low temperatures create conditions that can preserve tissue for centuries, possibly including the neurological basis of the human mind. Through a process called vitrification, brain tissue can be cooled to cryogenic temperatures without ice formation. Damage associated with this process is theoretically reversible in the same sense that rejuvenation is theoretically possible by specific foreseeable technology. Injury to the brain due to stopped blood flow is now known to result from a complex series of processes that take much longer to run to completion than the six minute limit of ordinary resuscitation technology. Reperfusion beyond the six minute limit primarily damages blood vessels rather than brain tissue. Apoptosis of neurons takes many hours. This creates a window of opportunity between legal death and irretrievable loss of life for human and animal subjects to be cryopreserved with possibility of future resuscitation. Under ideal conditions, the time interval between onset of clinical death and beginning of cryonics procedures can be reduced to less than a minute, but much longer delays could also be compatible with ultimate survival. Although the evidence that cryonics may work is indirect, indirect evidence is essential in many areas of science. If complex changes due to aging are reversible at some future date, then similarly complex changes due to stopped blood flow and cryopreservation may also be reversible, with life-saving results for anyone with medical needs that exceed current capabilities.

*Cryonics Institute, Clinton Township, Michigan.

CRYONICS OVERVIEW

Cryonics is the practice of preserving humans and animals at cryogenic temperatures in the hope that future science can restore them to a healthy living condition as well as rejuvenate them. At present cryonics can only be performed after pronouncement of legal death of the cryonics subject.

The scientific justification for the practice of cryonics is based on several key concepts: (1) Low temperature can slow metabolism. Sufficiently low temperature can virtually stop chemical changes for centuries. (2) Ice formation can be reduced or even eliminated by the use of vitrification mixtures. (3) Legally dead does not mean "irreversibly dead". Death is a process, not an event -- and the process takes longer than is commonly believed. (4) Damage associated with low temperature preservation and clinical death that is not reversible today is theoretically reversible in the future.

Pronouncement of legal death is necessary before cryonics procedures can begin because cryonics is not yet a proven, recognized medical procedure. Following legal death, a cryonics team can then begin preservation procedures immediately. Cryonics preservation procedures are intended to protect the tissues of cryonics subjects while cooling them to temperatures below -120°C with minimal alteration of tissue structure after cardiac arrest.

In the first stages the circulation and respiration of the cryonics subject is mechanically restored, the subject is administered protective medicines and is rapidly cooled to a temperature between 10°C and 0°C . The subject's blood is washed out and a significant amount of body water is replaced with a cryoprotectant mixture to prevent ice formation. The subject is cooled to a temperature below -120°C and held in cryostasis. When and if future medicine has the capability, the subject will be re-warmed, cryoprotectant will be removed, tissues will be repaired, diseases will be cured, and the subject will be rejuvenated (if required).

COOLING

Preservation of food in refrigerators and freezers is based on the principle of lowering temperature to reduce the rate of biochemical degradation. Cooling to reduce metabolic rate (and ultimately to bring chemical processes to a virtual halt) is at the heart of cryonics practice. Initial cooling after pronouncement of death involves placing the cryonics subject in a bath of ice water. Cardiopulmonary support with mechanical active compression/decompression also speeds cooling because of heat transfer from flowing blood.

Cryonics subjects are cooled with convection, a combination of conduction and fluid motion. In convection, a solid object (such as a cryonics subject) is cooled by a fluid (liquid or gas) that is rapidly circulated, such that the fluid can carry heat away from the conduction layer around the solid object. In cooling a cryonics subject from human body temperature (37°C) to 10°C , cooling by rapid circulation of ice-water is far more effective because of the convection effect than cooling by ice-packs or by standing water.

The formula governing convection is **Newton's Law of Cooling** which equates the rate of heat transfer to $hA(T_s - T_f)$ where T_s is the starting temperature (head or body of a cryonics subject), T_f is the final temperature (temperature of the cooling medium, e.g., ice water or cold nitrogen gas), A is the surface area of the solid, and h is a variable which is

dependent upon the rate of fluid motion as well as the thermal conductivity and heat capacity of the cooling medium. Faster fluid motion and higher thermal conductivity will increase the value of h . Newton's Law of Cooling predicts that cooling rate is greatest at the start of cooling when T_s is much greater than T_f . The cooling rate declines exponentially thereafter.

Reduction in temperature can considerably extend the time without blood flow before irreversible damage occurs. Many people, especially children, have been reported to survive 20 minutes to an hour or more of cardiac arrest with complete neurological recovery after hypothermic accidents, such as drowning in cold water^{1,2}. Metabolic rate can be dramatically reduced by cooling.

Duration of ischemic time necessary to cause 50% neuronal damage in gerbils has been shown to increase exponentially with lowering of brain temperature from 37°C to 31°C³. Six out of six experimental hypothermic dogs having tympanic temperature of 10°C were shown to endure 90 minutes of cardiac arrest without subsequent neurological damage, and two out of seven endured 120 minutes without evident neurological damage⁴. Humans have been subjected to deep hypothermic cardiac arrest for aortic surgery for over an hour without gross neurological deficits. The subjects reached complete electrocerebral silence (zero electroencephalographic bispectral index) between temperatures of 16°C and 24°C⁵ and were re-warmed without neurological deficit, confirming that dynamic brain activity can be lost and regained without loss of personal identity.

The extension of hypothermic protection from ischemic injury to subzero temperatures is seen in the northern wood frog (*Rana sylvatica*) which can survive in a semi-frozen state without heartbeat for months at temperatures as low as -3°C to -6°C with full recovery upon re-warming⁶. In 1966 a Japanese researcher replaced blood with glycerol to reduce ice formation in cat brains cooled to -20°C. After 45 days with no blood circulation at -20°C the revived cat brains demonstrated normal-looking EEG activity⁷.

The relationship between reaction rate (k) of chemical reactions (including metabolism and the processes of ischemic injury) and temperature (T) can be described by the Arrhenius equation⁸:

$$k = A \exp(-E_a/RT)$$

where T is in Kelvins, E_a is the **activation energy**, R is the **universal gas constant** (8.314 Joules/mole-Kelvin) and A is the **frequency factor** (related to frequency of molecular collisions and the probability that collisions are favorably oriented for reaction). Taking the natural logarithm of both sides of this equation gives:

$$\ln k = (- E_a/RT) + \ln A$$

For each of two different temperatures, T_1 and T_2 , there will be a different temperature-dependent reaction rate, k_1 and k_2 :

$$\ln k_1 = (- E_a/RT_1) + \ln A$$

$$\ln k_2 = (- E_a/RT_2) + \ln A$$

Subtracting $\ln k_2$ from $\ln k_1$ gives a single equation for the four variables:

$$\ln k_1 - \ln k_2 = ((- E_a/RT_1) + \ln A) - ((- E_a/RT_2) + \ln A)$$

which can be simplified to:

$$\ln (k_1/k_2) = (E_a/R)*(1/T_2 - 1/T_1)$$

or $k_1/k_2 = e^{(E_a/R)*(1/T_2 - 1/T_1)}$

The reaction rates of enzymes at various temperatures give a close approximation to the relationship between temperature and metabolic rate. Lactate dehydrogenase from rabbit muscle -- which has an activation energy (E_a) of 13,100 calories/mole⁹ -- can be taken as a representative enzyme. Using one thermochemical calorie equal to 4.184 Joules gives 54,810 Joules/mole.

Comparing the reaction rate (k_1) for lactate dehydrogenase at 40°C (313 Kelvins) (T_1) to the reaction rate (k_2) at 30°C (303 Kelvins) (T_2) gives:

$$k_1/k_2 = e^{((54,810 \text{ J/mol})/(8.314 \text{ J/mol-K})*(1/303 \text{ K} - 1/313 \text{ K}))} = 2.004$$

The reaction rate at 40°C is almost exactly twice the reaction rate at 30°C or, conversely, dropping the temperature 10°C has the effect of cutting the reaction rate nearly in half. This is in agreement with the **Q₁₀ rule**, a rule of thumb that between 0°C and 40°C reaction rates are reduced by one-half to one-third for every 10°C drop in temperature¹⁰.

This exponential drop in reaction rates with declining temperature means that reaction rates would become infinitesimally small at cryogenic temperatures (temperatures below -100°C) if chemical reactions were possible at those temperatures. The following table, produced by using the previous equation, compares the reaction rate at 37°C (310 Kelvins, normal human body temperature) to reaction rates at lower temperatures.

Reaction rate at 37°C compared to lower temperatures

Temperature	Reference	Relative rate at 37°C	Relative to 6 min. at 37°C
0°C (273K)	melting ice	18	1.8 hours
-80°C (193K)	dry ice	400,000	4.5 years
-120°C (153K)	glass transition	3 billion (3*10 ⁹)	34,000 years
-196°C (77K)	boiling nitrogen	9 octillion (9*10 ²⁷)	100 sextillion (10 ²³) years

If lactate dehydrogenase reaction rate was representative of metabolism in general, the metabolism at 37°C would be 18 times faster than at 0°C. Experimentally it has been

observed that the rate of oxidative phosphorylation at 4°C is about one-twentieth the rate at 37°C¹¹, a figure roughly in agreement with the value just calculated.

A reaction rate that is 9 octillion times faster at human body temperature than at -196°C would indicate essentially no reaction for millennia at the lower temperature. At this rate it would take 100 sextillion years for the ischemic biochemical reactions that occur at 37°C in six minutes to occur at liquid nitrogen temperature. But even these figures understate chemical inertness at lower cryogenic temperatures because the Arrhenius equation is based on the assumption of a fluid or gas medium in which normal chemistry is possible. Below -130°C even vitrified mammalian tissues are in a solid state, with a viscosity in excess of 10¹³ poise^{12,13}, a viscosity about 10¹⁵ (one quadrillion) times greater than the viscosity of water at 20°C¹⁴. The resulting diffusion rates are insignificant over geological time spans. At liquid nitrogen temperature mammalian tissues would even be stable against background radiation over periods of many centuries¹².

It is a misconception that freezing mammalian tissue typically results in ice formation within cells, causing the cells to burst. As mammalian tissues are cooled water leaves cells osmotically to form extracellular pure water-ice crystals. The unfrozen solution will contain increasing concentrations of toxic electrolytes. Ultimately enough extracellular ice will form to crush cells in the remaining unfrozen channels¹². Whether mechanical crushing or toxic electrolytes is the cause of damage following ice formation during slow cooling remains a subject of debate among cryobiologists¹⁵. Cryonics practice is based on efforts to reduce or eliminate freezing, however.

VITRIFICATION AND CRYOGENIC STORAGE

Cryonics practice has long sought to minimize ice formation by perfusing cryonics subjects with anti-freeze compounds known as cryoprotectants, traditionally glycerol. As of 2007 both of the major cryonics organizations doing cryoprotectant perfusions (Alcor Life Extension Foundation and the Cryonics Institute) claim to have eliminated ice formation in the brain by the use of vitrification solution, but make no such claim for other organs or tissues^{16,17}.

Vitrification is solidification to an amorphous (glassy) state which is distinct from the crystalline state characteristic of ice. Amber is a familiar example of a vitreous (amorphous, non-crystalline) solid. Pure water can be made to vitrify if cooled not more slowly than three million Kelvins per second¹⁸, a cooling rate impractical for animal tissues. Sucrose can be cooled rapidly enough to be vitrified into "cotton candy", but with slower cooling will form a "rock candy" crystal. Adding corn syrup to sucrose allows it to be cooled slowly to the non-crystalline solid used in lollipops. Silicon dioxide can be rapidly cooled to vitreous silica or can be slowly cooled to the crystalline form (quartz). Common glassware and window panes are made by adding sodium and calcium oxides to silicon dioxide to produce a molten liquid that can cool slowly as an increasingly viscous syrup to an amorphous (non-crystalline) solid. In the absence of a phase transition from liquid to solid crystal at melting/fusion temperature, there is a great increase in viscosity

(characterized as solidification) which occurs at a glass transition temperature (T_g) that is determined by cooling rate.

Cryoprotectants most frequently used in cryobiology include dimethylsulfoxide (DMSO) as well as the polyols ethylene glycol (an automobile anti-freeze), propylene glycol (once used to reduce ice crystals in ice cream) and glycerol (used since the 1950s to cryopreserve sperm and blood cells). All of these compounds are capable of hydrogen bonding with water to prevent water molecules from organizing themselves into ice. These cryoprotectants also act by colligative interference that hinders water molecules from forming the ice lattice. Mixtures of cryoprotectants can be less toxic than the pure cryoprotectants, and can completely eliminate ice formation. The use of ice blockers (non-cryoprotectant substances such as anti-freeze proteins that chemically block ice crystal growth) in vitrification mixtures can further reduce toxicity and concentration needed to vitrify¹⁹.

Difficulty in achieving sufficiently high cryoprotectant concentration to eliminate ice formation, while at the same time minimizing cryoprotectant toxicity, has been the limiting factor preventing better recovery of biological systems from cryopreservation. Rapid cooling can permit the use of lower cryoprotectant concentrations to prevent ice formation, but rapid cooling becomes increasingly difficult for increasingly larger tissues. Cryoprotectant toxicity varies inversely with temperature, so the use of less viscous cryoprotectant mixtures can speed tissue penetration and thereby reduce the tissue cryoprotectant exposure time at higher temperatures before cooling.

A number of possible explanations for cryoprotectant toxicity have been proposed, but the exact molecular mechanisms remain elusive²⁰. Insofar as cryoprotectants do not destroy molecules, the damage they cause may not be irreparable. Moreover, considerable success has been made in reducing the toxicity of vitrification mixtures^{21,22}, and there is no reason to believe that further toxicity reductions cannot be made.

The mammalian organ which has been studied by the most researchers attempting organ vitrification is the ovary. Variable success has been achieved with the ovaries of a number of species, but the greatest success has been with the mouse ovary. Vitrified mouse ovaries cryopreserved at -196°C have been re-warmed to produce live-pup birth rates comparable to that seen with fresh ovaries²³.

A study on rat hippocampal slices showed that it is possible for vitrified slices cooled to a solid state at -130°C to have viability upon re-warming comparable to that of control slices that had not been vitrified or cryopreserved. Ultrastructure of the CA1 region (the region of the brain most vulnerable to ischemic damage) of the re-warmed slices is seen to be quite well preserved compared to the ultrastructure of control CA1 tissue²⁴. Cryonics organizations perfuse brains with vitrification solution until saturation is achieved.

Tissues which have been vitrified and cryopreserved are assessed for viability as well as for ultrastructure. Intracellular K^+/Na^+ ratio is a commonly used method of assessing

viability, although other methods (such as measurement of intracellular ATP content) could be useful in the future. The sodium pump which maintains membrane potential will not function without binding to ATP and Na^+ inside the membrane and K^+ outside the membrane. Although a cell can maintain a membrane potential for several hours without a functional sodium pump, the slow leak of Na^+ into the cell and consequent leak of K^+ out of the cell will result in a complete loss of membrane potential after several hours. Similarly, if the cell dies in the sense of no longer being capable of producing energy (ATP) in the mitochondria, the sodium pump will cease to operate. Thus, normal intracellular K^+/Na^+ ratios indicate functioning sodium pumps and intact cell membranes.

To assay the intracellular K^+/Na^+ ratio tissues are placed in mannitol to wash away extracellular ions. Then Trichloroacetic acid is used to rupture cell membranes and release intracellular ions. A flame photometer or atomic absorption spectrometer can be used to determine the relative concentrations of sodium and potassium ions. Viability studies of vitrified hippocampal slices using intracellular K^+/Na^+ ratios indicated viability in excess of 90% normal²⁴.

A rabbit kidney has been vitrified, cooled to -135°C , re-warmed and transplanted into a rabbit. The formerly vitrified transplant functioned well enough as the sole kidney to keep the rabbit alive indefinitely²⁵. Some people imagine a need to understand brain function as being essential for brain cryopreservation. But in simple terms, a kidney produces urine and a brain produces consciousness. A re-warmed brain that is physiologically restored should be able to produce consciousness no less than a re-warmed vitrified kidney can produce urine. Preservation of structure and restoration of physiology should result in restoration of function, irrespective of the organ or tissue.

The vitrification mixture used in preserving the rabbit kidney is known as M22. M22 is used by the cryonics organization Alcor for vitrifying cryonics subjects. Perfusion of rabbits with M22 has been shown to preserve brain ultrastructure without ice formation²⁶.

Cooling from 0°C to -130°C should be rapid to minimize the possibility of ice formation. When cooling from -130°C to -196°C thermal stress on large solid vitrified samples can cause cracking and fracturing²⁷. Although it should theoretically be possible to cool to -196°C slowly enough to avoid cracking, the requisite cooling rates are unknown and may be too slow to be practical. Annealing a vitrified sample near glass transition temperature can reduce thermal stress²⁸, but this may not be adequate. Due to its more well-defined nature, cracking damage may be much easier to repair than freezing damage.

"REVERSIBLE DEATH" ?

As recently as the 1950s it was believed that death is irreversible when the heart stops. Today it is established that CardioPulmonary Resuscitation (CPR) in combination with Automated External Defibrillators (AEDs) can restore many people to life who were clinically dead because of cardiac arrest²⁹. But it is still widely believed that after about

six minutes of cardiac arrest without circulation irreparable brain damage has already occurred.

In 1976 Peter Safar (the "father of CPR") showed that dogs could be subjected to twelve minutes of cardiac arrest without neurological damage by the use of elevated arterial pressure, norepinephrine, heparin, and hemodilution with dextran 40³⁰. Over a decade later an experiment showed that spontaneous EEG activity returned in 50% of cats subjected to one hour of global cerebral ischemia followed by reperfusion and treatment with norepinephrine (or dopamine), heparin, insulin, and acidosis buffers. Six out of fifteen of the cats submitted to intensive care regained spontaneous respiration, and one of those cats survived a full year with normal neurological function (except slight ataxia)³¹. The six minute limit is not mainly a neurological phenomenon, it is a problem of increased vascular resistance that can be overcome (in part) by increasing perfusion pressure³².

Reperfusion injury refers to the tissue damage inflicted when blood flow is restored after an ischemic period of more than about twenty minutes. The re-supply of blood after an excessive period of ischemia initiates inflammatory processes and causes oxygen to form toxic free radicals (reactive oxygen species) such as superoxide³³. Xanthine oxidase-produced superoxide damages the endothelium far more than the parenchyma³⁴. In inflammatory conditions, such as occur in reperfusion, inducible nitric oxide synthetase can increase nitric oxide concentration to thousands of times normal levels³⁵. During reperfusion, abnormally high amounts of superoxide convert almost all available nitric oxide to peroxynitrite -- regarded as the agent causing most of the damage to brain capillary endothelial cells³⁶.

Despite the damaging effects of excitotoxicity³⁷ brain structure is normally retained post-mortem much longer than is commonly appreciated. In the cerebral cortex of rats subjected to occlusion of blood flow to the cortex (cerebral ischemia) only 15% of neurons were necrotic after 6 hours. Most neurons (65%) did not become necrotic until 12 hours after the cessation of blood flow³⁸. Neurons isolated from the brains of autopsied elderly humans an average of 2.6 hours post-mortem showed 70-90% viability after two weeks *in vitro*³⁹.

Part of the reason why more than six minutes of cardiac arrest currently leads to neurological damage is because the ischemia starts a process of neuron self-destruction (apoptosis) which takes many hours to complete. But therapies are on the horizon which may interfere with apoptosis. Neurons in the CA1 sector of the hippocampus are much more vulnerable to becoming necrotic following ischemia than neurons elsewhere in the brain⁴⁰. But cell death in the hippocampus following ischemia can be significantly reduced by the use of caspase inhibitors that arrest the apoptotic process⁴¹. Caspase inhibitors have also been used to block apoptosis in cryopreserved hematopoietic cells re-warmed from cryogenic temperatures⁴². Bag-1 protein which binds pro-apoptotic members of the Bcl-2 protein family has demonstrated powerful anti-apoptotic effects on rat livers subjected to ischemia/reperfusion injury⁴³.

Most neuroscientists agree that the anatomical basis of mind is encoded in physical structures of the brain, particularly neuropil connectivity and synaptic strengths⁴⁴ and possibly neuronal epigenetic structure⁴⁵. The fact that complete absence of electrical activity in the brain does not prevent full neurological recovery^{5,46} supports the proposition that the ultimate basis of consciousness is structural rather than dynamic and can therefore be preserved at cryogenic temperatures.

Possibly associated with redundancy of information storage in the brain, significant recovery of cerebral cortex function is possible following stroke^{47,48,49}. Neural stem cell transplantation therapies have the potential to further augment brain recovery from damage due to ischemia, toxins and cryopreservation injury⁵⁰. These considerations increase the amount of damage that may be tolerable for restoration of a human cryonically preserved under suboptimal conditions.

Preservation of brain structure and restoration of brain function are essential to cryonics. Other organs and tissues are not as important because artificial organs and tissue regeneration by stem cells should be easily accomplished by future medicine. Appendage regeneration in salamanders is already being used as a guideline for mammalian regenerative medicine⁵¹. Biodegradable 3-dimensional braided fibrous scaffolds⁵² can potentially be used for construction of organs, if not whole bodies.

Conservative cryonics strives to minimize damage and minimize reliance on future molecular repair technologies. In many cases cryonics subjects have experienced less than a minute of cardiac arrest before circulation has been restored. Evidence that the neurological basis of mind is preserved long beyond the six minute limit gives hope that molecular medicine to reverse apoptosis and repair damaged blood vessels may allow for recovery of cryonics subjects who did not benefit from prompt treatment. It is unlikely that cryonics is worthless after six minutes without blood circulation. Many tissues are alive when the heart stops, and take hours to die.

Under the best circumstances, cryonics subjects experience virtually no ice formation in the brain. Repairs to vitrified brain tissue that had experienced little ischemic damage could be performed above cryogenic temperatures along with curing diseases and rejuvenating.

Although there are sound legal requirements for drawing a distinct line between life and death, biological and psychological reality point to a continuum rather than discrete binary states. Consciousness emerges gradually from embryo to fetus to child to adult, and consciousness can diminish gradually in neurodegenerative diseases. After the heart stops the anatomical basis of mind (the brain) decomposes over a period of hours and days at a temperature-dependent rate. The non-binary character of consciousness would also become evident in revived cryonics subjects whose brains had been partially destroyed and then repaired, resulting in partial amnesia and partial restoration of original identity.

CRYONICS PROCEDURES

Pre-treatment of terminal cryonics subjects to reduce ischemia/reperfusion injury is advisable, but is too rarely done in practice. For example, intravenous injection of the alpha-tocopherol form of Vitamin E (20 mg/kg) 30 minutes prior to ischemia has been shown to significantly reduce lipid peroxidation and neurological damage⁵³. It is better to include both alpha-tocopherol and gamma-tocopherol because gamma-tocopherol removes peroxynitrite whereas alpha-tocopherol does not⁵⁴. Vitamin E pre-treatment for cryonics patients has the additional advantage of reducing blood clotting -- and does not have the risk of gastric bleeding associated with aspirin. Many fish oils (especially salmon oil) afford the same benefit, in addition to reducing the risk of cardiac arrest⁵⁵. Reduced clotting in a cryonics patient is usually a great benefit. But for patients undergoing surgery, Vitamin E and fish oils may be prohibited because of the danger of excessive bleeding.

Cryonics procedures are generally only practiced on subjects who have made contractual and funding arrangements in advance with a cryonics organization (such as Alcor Life Extension Foundation, the American Cryonics Society or the Cryonics Institute). In optimal circumstances a cryonics subject will be pronounced legally dead very quickly after their heart stops. Only after legal death has been pronounced can the cryonics procedures begin.

Once cardiac arrest has occurred and death has been pronounced, a cryonics subject can be given medications to maintain sedation, reduce cerebral metabolism, prevent/reverse blood clotting, increase blood pressure, stabilize pH against acidosis, and protect against ischemia/reperfusion injury.

Cryonics procedures involve restoring blood circulation and respiration as soon as possible to keep tissues alive. In cryonics, this is called CardioPulmonary Support (CPS) rather than CardioPulmonary Resuscitation (CPR) because resuscitation after death has been pronounced is not desired (a DNR, Do Not Resuscitate, condition). Propofol (2,6-diisopropylphenol) is given partly because its sedative action can prevent resuscitation, with the added benefit that it can be neuroprotective⁵⁶. Propofol has been shown to inhibit the neural cell apoptosis that can occur as a consequence of ischemia/reperfusion injury⁵⁷.

Heparin is used to prevent blood clotting. Streptokinase is the usual thrombolytic used to break up blood clots. THAM (Tris-Hydroxymethyl AminoMethane) is a buffer that maintains arterial pH without producing carbon dioxide and also maintains intracellular pH because it readily crosses cell membranes⁵⁸.

When the equipment is available, cryonics teams restore circulation and respiration with mechanical devices capable of restoring circulation on the down-stroke (compression) as well as the up-stroke (decompression). Active Compression-DeCompression (ACDC) and interposed abdominal compression can improve CPS perfusion considerably⁵⁹. Epinephrine has commonly been used to supplement CPS by maintaining blood pressure, although vasopressin may also be used⁶⁰.

While the cryonics subject is receiving ACDC CPS, he or she is in a bath of circulating ice water. Cooling is much more rapid in water than in air⁶¹ and is much more rapid in flowing water than in still water due to Newton's Law of Cooling. The cooling of the cryonics subject is also considerably hastened by the blood circulation resulting from ACDC CPS.

Once the cryonics subject is cooled to below 10°C perfusion with vitrification solution can begin. Vitrification of the brain is achieved by cryoprotectant exposure times which are considerably longer than was seen for vitrification of hippocampal slices. Cryoprotectants are toxic, so increased cryoprotectant exposure time means increased toxicity to the exposed tissues. But large organs and body tissues cannot be cooled as rapidly as tissue slices, therefore higher cryoprotectant concentrations must be used to prevent ice formation.

Suggestions have been made for systems to store cryonics subjects at temperatures closer to -130°C to eliminate cracking due to thermal stress in cooling to -196°C. Subjects stored just below -130°C would still be in the solid state, insofar as vitrification solutions have a glass transition temperature just below -120°C²⁰. Storage just below -130°C has yet to be implemented on all but a few cryonics subjects.

Cryonics subjects are stored in thermos-bottle-like containers of liquid nitrogen, a storage method which is both inexpensive and not dependent upon electricity (not so vulnerable to power failure).

SCIENCE AND INDIRECT EVIDENCE

Many critics assert that cryonics is not science and has no capability of becoming science until a mammal has been re-warmed and reanimated after having been cryopreserved at cryogenic temperatures. But model-building based on extrapolations from indirect evidence is central to science.

No one has ever seen the core of the Earth. Models have been constructed of the state of the universe in the first millionth of a second following the Big Bang. Scientists describe the state of the Earth after years of global warming. Models have been constructed of the state of contained nuclear waste hundreds of thousands of years in the future -- models upon which considerable reliance is placed in the disposal of nuclear waste. Past landings by humans on the moon provide indirect evidence that future landings of humans on Mars may be possible.

Many people cryopreserve umbilical cord stem cells from their newborn baby on the basis of potential future scientific developments rather than on the basis of current science. Germ cells and DNA from endangered species are similarly cryopreserved in anticipation of future technology. Influenza vaccine only has a 50% chance of protecting a person over 65^{62,63}. It is not unscientific to risk modest or heroic medical treatments that are justified by indirect evidence for some probability of success, rather than absolute guarantee of success.

If there are plausible models for the repair and reanimation of cryopreserved cryonics subjects^{64,65} it seems reasonable to rely upon them when deciding upon human cryopreservation as a long-term treatment which may or may not succeed. To wait until a cryopreserved mammal has been reanimated before cryopreserving humans could mean that many lives will be lost. It would be like waiting tens or hundreds of thousands of years to ensure that nuclear waste can be contained before containing those wastes. Indirect evidence exists to support the claim that revival of a cryopreserved mammal is not essential for cryonics practice to be scientifically justified.

CONCLUSIONS

Low temperatures slow biological time, effectively stopping time at liquid nitrogen temperature. Cryoprotectants greatly reduce damage caused by tissue cryopreservation, and effective vitrifications can prevent ice formation completely. Cryoprotectant toxicity is likely a repairable injury, and less toxic means of cryopreservation continue to be discovered. Applied to humans and animals, cryopreservation is a means of achieving a stable biological state that is reversible in principle.

Dying is a process that begins, not ends, when heartbeat and blood circulation stop. When legal death is declared based on cardiopulmonary arrest, cryonics procedures aim to minimize further injury by artificially restoring blood circulation and rapidly reducing temperature. The duration of clinical death at warm temperatures beyond which the brain information that defines a human being is lost may be many hours.

The proposition that aging is a disease that can be treated and perhaps eventually be reversed (rejuvenation), is based on the general understanding that aging consists of a multitude of specific pathologies on cellular and molecular levels that can be studied, understood, and reversed with foreseeable tools. Pathologies caused by global cerebral ischemia (clinical death), by cryopreservation, and those of other presently-incurable diseases are similarly amenable to analysis and possible future repair. If aging damage can be repaired at some future time, it is not unreasonable to think that damage due to cryonics procedures can also be repaired. And if aging damage can be repaired at some future time, cryonics may be the only way for many people living today to reach future medicine that can cure presently-incurable diseases and rejuvenate.

REFERENCES

1. Eich C, Brauer A, Kettler D. Recovery of a hypothermic drowned child after resuscitation with cardiopulmonary bypass followed by prolonged extracorporeal membrane oxygenation. *Resuscitation*. 2005 Oct;67(1):145-8.
2. Bolte RG, Black PG, Bowers RS, Thorne JK, Corneli HM. The use of extracorporeal rewarming in a child submerged for 66 minutes. *JAMA*. 1988 Jul 15;260(3):377-9.

3. Takeda Y, Namba K, Higuchi T, Hagioka S, Takata K, Hirakawa M, Morita K. Quantitative evaluation of the neuroprotective effects of hypothermia ranging from 34 degrees C to 31 degrees C on brain ischemia in gerbils and determination of the mechanism of neuroprotection. *Crit Care Med.* 2003 Jan;31(1):255-60.
4. Behringer W, Safar P, Wu X, Kentner R, Radovsky A, Kochanek PM, Dixon CE, Tisherman SA. Survival without brain damage after clinical death of 60-120 mins in dogs using suspended animation by profound hypothermia. *Crit Care Med.* 2003 May;31(5):1523-31.
5. Hayashida M, Sekiyama H, Orii R, Chinzei M, Ogawa M, Arita H, Hanaoka K, Takamoto S. Effects of deep hypothermic circulatory arrest with retrograde cerebral perfusion on electroencephalographic bispectral index and suppression ratio. *J Cardiothorac Vasc Anesth.* 2007 Feb;21(1):61-7.
6. Costanzo JP, Lee RE Jr, DeVries AL, Wang T, Layne JR Jr. Survival mechanisms of vertebrate ectotherms at subfreezing temperatures: applications in cryomedicine. *FASEB J.* 1995 Mar;9(5):351-8.
7. Suda I, Kito K, Adachi C. Viability of long term frozen cat brain in vitro. *Nature.* 1966 Oct 15;212(5059):268-70.
8. CHEMISTRY: The Central Science (Seventh Edition); pp.511-512; Brown, TL, et. al., Editors; Prentice Hall (1997).
9. Low PS, Bada JL, Somero GN. Temperature adaptation of enzymes: roles of the free energy, the enthalpy, and the entropy of activation. *Proc Natl Acad Sci U S A.* 1973 Feb;70(2):430-2.
10. Davidson EA, Janssens IA. Temperature sensitivity of soil carbon decomposition and feedbacks to climate change. *Nature.* 2006 Mar 9;440(7081):165-73.
11. Dufour S, Rousse N, Canioni P, Diolez P. Top-down control analysis of temperature effect on oxidative phosphorylation. *Biochem J.* 1996 Mar 15;314 (Pt 3):743-51.
12. Mazur P. Freezing of living cells: mechanisms and implications. *Am J Physiol.* 1984 Sep;247(3 Pt 1):C125-42.
13. Angell CA. Liquid fragility and the glass transition in water and aqueous solutions. *Chem Rev.* 2002 Aug;102(8):2627-50.
14. CRC Handbook of Chemistry and Physics (76th Edition), p.6-10, David R. Lido, Editor-in-Chief CRC Press (1995-1996).

15. Mazur,P, "Principles of Cryobiology" in *Life In the Frozen State*, pp.37-51, Barry J. Fuller, et. al., Editors; CRC Press (2004).
16. Chamberlain,F, *Vitrification arrives!* *Cryonics* 2000 21(4):4-9.
17. Best,B, *Long Life: Longevity through Technology*, CI-VM-1 cryoprotectant and CI carrier solution used for vitrification 2007 Mar-Apr; 39(3-4):5-6 .
18. Bald WB. *On crystal size and cooling rate.* *J Microsc.* 1986 Jul;143(Pt 1):89-102.
19. Wowk B, Leitl E, Rasch CM, Mesbah-Karimi N, Harris SB, Fahy GM. *Vitrification enhancement by synthetic ice blocking agents.* *Cryobiology.* 2000 May;40(3):228-36.
20. Fahy GM, Lilley TH, Linsdell H, Douglas MS, Meryman HT. *Cryoprotectant toxicity and cryoprotectant toxicity reduction: in search of molecular mechanisms.* *Cryobiology.* 1990 Jun;27(3):247-68.
21. Fahy GM, Wowk B, Wu J, Paynter S. *Improved vitrification solutions based on the predictability of vitrification solution toxicity.* *Cryobiology.* 2004 Feb;48(1):22-35.
22. Fahy GM, Wowk B, Wu J, Phan J, Rasch C, Chang A, Zendejas E. *Cryopreservation of organs by vitrification: perspectives and recent advances.* *Cryobiology.* 2004 Apr;48(2):157-78.
23. Hasegawa A, Mochida N, Ogasawara T, Koyama K. *Pup birth from mouse oocytes in preantral follicles derived from vitrified and warmed ovaries followed by in vitro growth, in vitro maturation, and in vitro fertilization.* *Fertil Steril.* 2006 Oct;86 Suppl 4:1182-92.
24. Pichugin Y, Fahy GM, Morin R. *Cryopreservation of rat hippocampal slices by vitrification.* *Cryobiology.* 2006 Apr;52(2):228-40.
25. Fahy,GM, "Vitrification as an approach to cryopreservation: general perspectives", *CRYOBIOLOGY*; 51(3):348 (2005) [Abstract].
26. Lemler J, Harris SB, Platt C, Huffman TM. *The arrest of biological time as a bridge to engineered negligible senescence.* *Ann N Y Acad Sci.* 2004 Jun;1019:559-63.
27. Fahy GM, Saur J, Williams RJ. *Physical problems with the vitrification of large biological systems.* *Cryobiology.* 1990 Oct;27(5):492-510.
28. Baudot,A., Odagescu,V, "An automated cryostat designed to study the vitrification of cryoprotective solutions without any function", *CRYOBIOLOGY* 53:442 (2006) [Abstract].

29. Cobb LA, Fahrenbruch CE, Walsh TR, Copass MK, Olsufka M, Breskin M, Hallstrom AP. Influence of cardiopulmonary resuscitation prior to defibrillation in patients with out-of-hospital ventricular fibrillation. *JAMA*. 1999 Apr 7;281(13):1182-8.
30. Safar P, Stezoski W, Nemoto EM. Amelioration of brain damage after 12 minutes' cardiac arrest in dogs. *Arch Neurol*. 1976 Feb;33(2):91-5.
31. Hossmann KA. Resuscitation potentials after prolonged global cerebral ischemia in cats. *Crit Care Med*. 1988 Oct;16(10):964-71.
32. Shaffner DH, Eleff SM, Brambrink AM, Sugimoto H, Izuta M, Koehler RC, Traystman RJ. Effect of arrest time and cerebral perfusion pressure during cardiopulmonary resuscitation on cerebral blood flow, metabolism, adenosine triphosphate recovery, and pH in dogs. *Crit Care Med*. 1999 Jul;27(7):1335-42.
33. de Groot H, Rauen U. Ischemia-reperfusion injury: processes in pathogenetic networks: a review. *Transplant Proc*. 2007 Mar;39(2):481-4.
34. Ratych RE, Chuknyiska RS, Bulkley GB. The primary localization of free radical generation after anoxia/reoxygenation in isolated endothelial cells. *Surgery*. 1987 Aug;102(2):122-31.
35. Brown GC, Borutaite V. Nitric oxide inhibition of mitochondrial respiration and its role in cell death. *Free Radic Biol Med*. 2002 Dec 1;33(11):1440-50.
36. Wu S, Tamaki N, Nagashima T, Yamaguchi M. Reactive oxygen species in reoxygenation injury of rat brain capillary endothelial cells. *Neurosurgery*. 1998 Sep;43(3):577-83.
37. Lipton P. Ischemic cell death in brain neurons. *Physiol Rev*. 1999 Oct;79(4):1431-568.
38. Garcia JH, Liu KF, Ho KL. Neuronal necrosis after middle cerebral artery occlusion in Wistar rats progresses at different time intervals in the caudoputamen and the cortex. *Stroke*. 1995 Apr;26(4):636-42.
39. Konishi Y, Lindholm K, Yang LB, Li R, Shen Y. Isolation of living neurons from human elderly brains using the immunomagnetic sorting DNA-linker system. *Am J Pathol*. 2002 Nov;161(5):1567-76.
40. Radovsky A, Safar P, Sterz F, Leonov Y, Reich H, Kuboyama K. Regional prevalence and distribution of ischemic neurons in dog brains 96 hours after cardiac arrest of 0 to 20 minutes. *Stroke*. 1995 Nov;26(11):2127-33.
41. Chen J, Nagayama T, Jin K, Stetler RA, Zhu RL, Graham SH, Simon RP. Induction of caspase-3-like protease may mediate delayed neuronal death in the hippocampus after transient cerebral ischemia. *J Neurosci*. 1998 Jul 1;18(13):4914-28.

42. Stroh C, Cassens U, Samraj AK, Sibrowski W, Schulze-Osthoff K, Los M. The role of caspases in cryoinjury: caspase inhibition strongly improves the recovery of cryopreserved hematopoietic and other cells. *FASEB J.* 2002 Oct;16(12):1651-3.
43. Sawitzki B, Amersi F, Ritter T, Fisser M, Shen XD, Ke B, Busuttill R, Volk HD, Kupiec-Weglinski JW. Upregulation of Bag-1 by ex vivo gene transfer protects rat livers from ischemia/reperfusion injury. *Hum Gene Ther.* 2002 Aug 10;13(12):1495-504.
44. Abraham WC, Robins A. Memory retention--the synaptic stability versus plasticity dilemma. *Trends Neurosci.* 2005 Feb;28(2):73-8.
45. Arshavsky YI. "The seven sins" of the Hebbian synapse: can the hypothesis of synaptic plasticity explain long-term memory consolidation *Prog Neurobiol.* 2006 Oct;80(3):99-113.
46. Rothstein TL. Recovery from near death following cerebral anoxia: A case report demonstrating superiority of median somatosensory evoked potentials over EEG in predicting a favorable outcome after cardiopulmonary resuscitation. *Resuscitation.* 2004 Mar;60(3):335-41.
47. Dancause N, Barbay S, Frost SB, Plautz EJ, Chen D, Zoubina EV, Stowe AM, Nudo RJ. Extensive cortical rewiring after brain injury. *J Neurosci.* 2005 Nov 2;25(44):10167-79.
48. Carmichael ST. Cellular and molecular mechanisms of neural repair after stroke: making waves. *Ann Neurol.* 2006 May;59(5):735-42.
49. Nudo RJ. Postinfarct cortical plasticity and behavioral recovery. *Stroke.* 2007 Feb;38(2 Suppl):840-5.
50. Savitz SI, Dinsmore JH, Wechsler LR, Rosenbaum DM, Caplan LR. Cell therapy for stroke. *NeuroRx.* 2004 Oct;1(4):406-14.
51. Brookes JP, Kumar A.
Appendage regeneration in adult vertebrates and implications for regenerative medicine.
Science. 2005 Dec 23;310(5756):1919-23.
52. Cooper JA Jr, Sahota JS, Gorum WJ 2nd, Carter J, Doty SB, Laurencin CT.
Biomimetic tissue-engineered anterior cruciate ligament replacement.
Proc Natl Acad Sci U S A. 2007 Feb 27;104(9):3049-54.
53. Yamamoto M, Shima T, Uozumi T, Sogabe T, Yamada K, Kawasaki T. A possible role of lipid peroxidation in cellular damages caused by cerebral ischemia and the protective effect of alpha-tocopherol administration. *Stroke.* 1983 Nov-Dec;14(6):977-82.

54. Christen S, Woodall AA, Shigenaga MK, Southwell-Keely PT, Duncan MW, Ames BN. Gamma-tocopherol traps mutagenic electrophiles such as NO(X) and complements alpha-tocopherol: physiological implications. *Proc Natl Acad Sci U S A*. 1997 Apr 1;94(7):3217-22.
55. Charnock JS, McLennan PL, Abeywardena MY. Dietary modulation of lipid metabolism and mechanical performance of the heart. *Mol Cell Biochem*. 1992 Oct 21;116(1-2):19-25.
56. Adembri C, Venturi L, Tani A, Chiarugi A, Gramigni E, Cozzi A, Pancani T, De Gaudio RA, Pellegrini-Giampietro DE. Neuroprotective effects of propofol in models of cerebral ischemia: inhibition of mitochondrial swelling as a possible mechanism. *Anesthesiology*. 2006 Jan;104(1):80-9.
57. Polster BM, Basanez G, Young M, Suzuki M, Fiskum G. Inhibition of Bax-induced cytochrome c release from neural cell and brain mitochondria by dibucaine and propranolol. *J Neurosci*. 2003 Apr 1;23(7):2735-43.
58. Gehlbach BK, Schmidt GA. Bench-to-bedside review: treating acid-base abnormalities in the intensive care unit - the role of buffers. *Crit Care*. 2004 Aug;8(4):259-65.
59. Babbs CF. CPR techniques that combine chest and abdominal compression and decompression: hemodynamic insights from a spreadsheet model. *Circulation*. 1999 Nov 23;100(21):2146-52.
60. Wenzel V, Lindner KH. Arginine vasopressin during cardiopulmonary resuscitation: laboratory evidence, clinical experience and recommendations, and a view to the future. *Crit Care Med*. 2002 Apr;30(4 Suppl):S157-61.
61. Baccino E, Cattaneo C, Jouineau C, Poudoulec J, Martrille L. Cooling rates of the ear and brain in pig heads submerged in water: implications for postmortem interval estimation of cadavers found in still water. *Am J Forensic Med Pathol*. 2007 Mar;28(1):80-5.
62. Nichol KL, Nordin J, Mullooly J, Lask R, Fillbrandt K, Iwane M. Influenza vaccination and reduction in hospitalizations for cardiac disease and stroke among the elderly. *N Engl J Med*. 2003 Apr 3;348(14):1322-32.
63. Hak E, Buskens E, van Essen GA, de Bakker DH, Grobbee DE, Tacke MA, van Hout BA, Verheij TJ. Clinical effectiveness of influenza vaccination in persons younger than 65 years with high-risk medical conditions: the PRISMA study. *Arch Intern Med*. 2005 Feb 14;165(3):274-80.

64. Drexler KE. Molecular engineering: An approach to the development of general capabilities for molecular manipulation. Proc Natl Acad Sci U S A. 1981 Sep;78(9):5275-5278.

65. Freitas RA Jr. What is nanomedicine? Nanomedicine. 2005 Mar;1(1):2-9.